

CORRECTION

Open Access



Correction: Comparative evaluation of clinical and cerebrospinal fluid biomarker characteristics in rapidly and non-rapidly progressive Alzheimer's disease

Janne Marieke Herden^{1†}, Peter Hermann^{1*†}, Isabel Schmidt¹, Kathrin Dittmar¹, Sezgi Canaslan¹, Luise Weglage¹, Sabine Nuhn¹, Corinna Volpers¹, Astrid Schlung¹, Stefan Goebel^{1,2}, Fabian Kück³, Anna Villar-Piqué¹, Christian Schmidt^{1,4}, Dirk Wedekind⁵ and Inga Zerr^{1,2}

Correction: *Alzheimers Res Ther* 15, 106 (2023)
<https://doi.org/10.1186/s13195-023-01249-y>

Published online: 22 June 2023

Following publication of the original article [1], the authors identified an error in Fig. 2 (the color coding of APOE genotypes E2/2 and E3/4 is wrong). APOE2/E2 is displayed as most frequent genotype but in fact, it is the least frequent genotype. The correct figure is given below.

The original article [1] has been corrected.

Reference

1. Herden JM, Hermann P, Schmidt I, et al. Comparative evaluation of clinical and cerebrospinal fluid biomarker characteristics in rapidly and non-rapidly progressive Alzheimer's disease. *Alzheimers Res Ther.* 2023;15:106. <https://doi.org/10.1186/s13195-023-01249-y>.

[†]Janne Marieke Herden and Peter Hermann contributed equally as first authors.

The original article can be found online at <https://doi.org/10.1186/s13195-023-01249-y>.

*Correspondence:

Peter Hermann
peter.hermann@med.uni-goettingen.de

¹ Department of Neurology, Clinical Dementia Center and National Reference Center for CJD Surveillance, University Medical Center, Robert-Koch-Straße 40, 37075 Göttingen, Germany

² German Center for Neurodegenerative Diseases (DZNE), Göttingen, Germany

³ Department of Medical Statistics, University Medical Center Göttingen, Humboldtallee 32, 37073 Göttingen, Germany

⁴ Neurologische Gemeinschaftspraxis Am Groner Tor, Göttingen, Germany

⁵ Department of Psychiatry and Psychotherapy, University Medical Center, Von-Siebold-Straße 5, 37075 Göttingen, Germany






Fig. 2 ApoE Genotype distribution at baseline. ApoE Genotype distribution in diagnostic groups. ApoE, apolipoprotein E; AD, Alzheimer's disease; rpAD, rapidly progressive Alzheimer's disease